DRUG FACTS

Purpose Active ingredient(s)

Use(s)

Warnings

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product

Stop use and ask a doctor if

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 fever gets worse or lasts more than 3 days
- · new symptoms occur

redness or swelling is present
 These could be signs of a serious condition.

If pregnant or breast-feeding

Keep out of reach of children

Directions

do not take more than un ected (see overdose warming)		
ı	adults	take 2 caplets every 8 hours with water
ı		swallow whole – do not crush, chew or dissolve
ı		do not take more than 6 caplets in 24 hours
ı		do not use for more than 10 days unless directed by a doctor
ı		
ı	under 18 years of age	ask a doctor
ı		

Other information

- store at 20-25°C (68-77°F). Avoid excessive heat 40°C (104°F).
- do not use if carton is opened or neck wrap or foil inner seal imprinted with "Safety Seal®" is broken

see end panel for lot number and expiration date

Drug Facts

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
 with other drugs containing acetaminophen
 3 or more alcoholic drinks every day while using this product

Inactive ingredients

carnauba wax, corn starch, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin